

MATH 304
Linear Algebra

Lecture 30:
The Gram-Schmidt process (continued).

Orthogonal sets

Let V be a vector space with an inner product.

Definition. Nonzero vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in V$ form an **orthogonal set** if they are orthogonal to each other: $\langle \mathbf{v}_i, \mathbf{v}_j \rangle = 0$ for $i \neq j$.

If, in addition, all vectors are of unit norm, $\|\mathbf{v}_i\| = 1$, then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ is called an **orthonormal set**.

Theorem Any orthogonal set is linearly independent.

Orthogonal projection

Let V be an inner product space.

Let $\mathbf{x}, \mathbf{v} \in V$, $\mathbf{v} \neq \mathbf{0}$. Then $\mathbf{p} = \frac{\langle \mathbf{x}, \mathbf{v} \rangle}{\langle \mathbf{v}, \mathbf{v} \rangle} \mathbf{v}$ is the

orthogonal projection of the vector \mathbf{x} onto the vector \mathbf{v} . That is, the remainder $\mathbf{o} = \mathbf{x} - \mathbf{p}$ is orthogonal to \mathbf{v} .

If $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an orthogonal set of vectors then

$$\mathbf{p} = \frac{\langle \mathbf{x}, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 + \frac{\langle \mathbf{x}, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 + \cdots + \frac{\langle \mathbf{x}, \mathbf{v}_n \rangle}{\langle \mathbf{v}_n, \mathbf{v}_n \rangle} \mathbf{v}_n$$

is the **orthogonal projection** of the vector \mathbf{x} onto the subspace spanned by $\mathbf{v}_1, \dots, \mathbf{v}_n$. That is, the remainder $\mathbf{o} = \mathbf{x} - \mathbf{p}$ is orthogonal to $\mathbf{v}_1, \dots, \mathbf{v}_n$.

The Gram-Schmidt orthogonalization process

Let V be a vector space with an inner product. Suppose $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ is a basis for V . Let

$$\mathbf{v}_1 = \mathbf{x}_1,$$


$$\mathbf{v}_2 = \mathbf{x}_2 - \frac{\langle \mathbf{x}_2, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1,$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \frac{\langle \mathbf{x}_3, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \frac{\langle \mathbf{x}_3, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2,$$

.....


$$\mathbf{v}_n = \mathbf{x}_n - \frac{\langle \mathbf{x}_n, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \dots - \frac{\langle \mathbf{x}_n, \mathbf{v}_{n-1} \rangle}{\langle \mathbf{v}_{n-1}, \mathbf{v}_{n-1} \rangle} \mathbf{v}_{n-1}.$$

Then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an orthogonal basis for V .



Properties of the Gram-Schmidt process:

- $\mathbf{v}_k = \mathbf{x}_k - (\alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1})$, $1 \leq k \leq n$;
- the span of $\mathbf{v}_1, \dots, \mathbf{v}_k$ is the same as the span of $\mathbf{x}_1, \dots, \mathbf{x}_k$;
- \mathbf{v}_k is orthogonal to $\mathbf{x}_1, \dots, \mathbf{x}_{k-1}$;
- $\mathbf{v}_k = \mathbf{x}_k - \mathbf{p}_k$, where \mathbf{p}_k is the orthogonal projection of the vector \mathbf{x}_k on the subspace spanned by $\mathbf{x}_1, \dots, \mathbf{x}_{k-1}$;
- $\|\mathbf{v}_k\|$ is the distance from \mathbf{x}_k to the subspace spanned by $\mathbf{x}_1, \dots, \mathbf{x}_{k-1}$.



Problem. Let Π be the plane in \mathbb{R}^3 spanned by vectors $\mathbf{x}_1 = (1, 2, 2)$ and $\mathbf{x}_2 = (-1, 0, 2)$.

(i) Find an orthonormal basis for Π .

(ii) Extend it to an orthonormal basis for \mathbb{R}^3 .

$\mathbf{x}_1, \mathbf{x}_2$ is a basis for the plane Π . We can extend it to a basis for \mathbb{R}^3 by adding one vector from the standard basis. For instance, vectors $\mathbf{x}_1, \mathbf{x}_2$, and $\mathbf{x}_3 = (0, 0, 1)$ form a basis for \mathbb{R}^3 because

$$\begin{vmatrix} 1 & 2 & 2 \\ -1 & 0 & 2 \\ 0 & 0 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 2 \\ -1 & 0 \end{vmatrix} = 2 \neq 0.$$

Using the Gram-Schmidt process, we orthogonalize the basis $\mathbf{x}_1 = (1, 2, 2)$, $\mathbf{x}_2 = (-1, 0, 2)$, $\mathbf{x}_3 = (0, 0, 1)$:

$$\mathbf{v}_1 = \mathbf{x}_1 = (1, 2, 2),$$

$$\begin{aligned}\mathbf{v}_2 &= \mathbf{x}_2 - \frac{\langle \mathbf{x}_2, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 = (-1, 0, 2) - \frac{3}{9}(1, 2, 2) \\ &= (-4/3, -2/3, 4/3),\end{aligned}$$

$$\begin{aligned}\mathbf{v}_3 &= \mathbf{x}_3 - \frac{\langle \mathbf{x}_3, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \frac{\langle \mathbf{x}_3, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 \\ &= (0, 0, 1) - \frac{2}{9}(1, 2, 2) - \frac{4/3}{4}(-4/3, -2/3, 4/3) \\ &= (2/9, -2/9, 1/9).\end{aligned}$$

Now $\mathbf{v}_1 = (1, 2, 2)$, $\mathbf{v}_2 = (-4/3, -2/3, 4/3)$,
 $\mathbf{v}_3 = (2/9, -2/9, 1/9)$ is an orthogonal basis for \mathbb{R}^3
while $\mathbf{v}_1, \mathbf{v}_2$ is an orthogonal basis for Π . It remains
to normalize these vectors.

$$\langle \mathbf{v}_1, \mathbf{v}_1 \rangle = 9 \implies \|\mathbf{v}_1\| = 3$$

$$\langle \mathbf{v}_2, \mathbf{v}_2 \rangle = 4 \implies \|\mathbf{v}_2\| = 2$$

$$\langle \mathbf{v}_3, \mathbf{v}_3 \rangle = 1/9 \implies \|\mathbf{v}_3\| = 1/3$$

$$\mathbf{w}_1 = \mathbf{v}_1 / \|\mathbf{v}_1\| = (1/3, 2/3, 2/3) = \frac{1}{3}(1, 2, 2),$$

$$\mathbf{w}_2 = \mathbf{v}_2 / \|\mathbf{v}_2\| = (-2/3, -1/3, 2/3) = \frac{1}{3}(-2, -1, 2),$$

$$\mathbf{w}_3 = \mathbf{v}_3 / \|\mathbf{v}_3\| = (2/3, -2/3, 1/3) = \frac{1}{3}(2, -2, 1).$$

$\mathbf{w}_1, \mathbf{w}_2$ is an orthonormal basis for Π .

$\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$ is an orthonormal basis for \mathbb{R}^3 .

Problem. Find the distance from the point $\mathbf{y} = (0, 0, 0, 1)$ to the subspace $\Pi \subset \mathbb{R}^4$ spanned by vectors $\mathbf{x}_1 = (1, -1, 1, -1)$, $\mathbf{x}_2 = (1, 1, 3, -1)$, and $\mathbf{x}_3 = (-3, 7, 1, 3)$.

Let us apply the Gram-Schmidt process to vectors $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{y}$. We should obtain an orthogonal system $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$. The desired distance will be $|\mathbf{v}_4|$.

$$\mathbf{x}_1 = (1, -1, 1, -1), \quad \mathbf{x}_2 = (1, 1, 3, -1), \\ \mathbf{x}_3 = (-3, 7, 1, 3), \quad \mathbf{y} = (0, 0, 0, 1).$$

$$\mathbf{v}_1 = \mathbf{x}_1 = (1, -1, 1, -1),$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \frac{\langle \mathbf{x}_2, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 = (1, 1, 3, -1) - \frac{4}{4}(1, -1, 1, -1) \\ = (0, 2, 2, 0),$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \frac{\langle \mathbf{x}_3, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \frac{\langle \mathbf{x}_3, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 \\ = (-3, 7, 1, 3) - \frac{-12}{4}(1, -1, 1, -1) - \frac{16}{8}(0, 2, 2, 0) \\ = (0, 0, 0, 0).$$

The Gram-Schmidt process can be used to check linear independence of vectors!

The vector \mathbf{x}_3 is a linear combination of \mathbf{x}_1 and \mathbf{x}_2 .

Π is a plane, not a 3-dimensional subspace.

We should orthogonalize vectors $\mathbf{x}_1, \mathbf{x}_2, \mathbf{y}$.

$$\begin{aligned}\mathbf{v}_4 &= \mathbf{y} - \frac{\langle \mathbf{y}, \mathbf{v}_1 \rangle}{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle} \mathbf{v}_1 - \frac{\langle \mathbf{y}, \mathbf{v}_2 \rangle}{\langle \mathbf{v}_2, \mathbf{v}_2 \rangle} \mathbf{v}_2 \\ &= (0, 0, 0, 1) - \frac{-1}{4}(1, -1, 1, -1) - \frac{0}{8}(0, 2, 2, 0) \\ &= (1/4, -1/4, 1/4, 3/4).\end{aligned}$$

$$|\mathbf{v}_4| = \left| \left(\frac{1}{4}, -\frac{1}{4}, \frac{1}{4}, \frac{3}{4} \right) \right| = \frac{1}{4} |(1, -1, 1, 3)| = \frac{\sqrt{12}}{4} = \frac{\sqrt{3}}{2}.$$

Problem. Find the distance from the point $\mathbf{z} = (0, 0, 1, 0)$ to the plane Π that passes through the point $\mathbf{x}_0 = (1, 0, 0, 0)$ and is parallel to the vectors $\mathbf{v}_1 = (1, -1, 1, -1)$ and $\mathbf{v}_2 = (0, 2, 2, 0)$.

The plane Π is not a subspace of \mathbb{R}^4 as it does not pass through the origin. Let $\Pi_0 = \text{Span}(\mathbf{v}_1, \mathbf{v}_2)$. Then $\Pi = \Pi_0 + \mathbf{x}_0$.

Hence the distance from the point \mathbf{z} to the plane Π is the same as the distance from the point $\mathbf{z} - \mathbf{x}_0$ to the plane Π_0 .

We shall apply the Gram-Schmidt process to vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{z} - \mathbf{x}_0$. This will yield an orthogonal system $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$. The desired distance will be $|\mathbf{w}_3|$.

$$\mathbf{v}_1 = (1, -1, 1, -1), \mathbf{v}_2 = (0, 2, 2, 0), \mathbf{z} - \mathbf{x}_0 = (-1, 0, 1, 0).$$

$$\mathbf{w}_1 = \mathbf{v}_1 = (1, -1, 1, -1),$$

$$\mathbf{w}_2 = \mathbf{v}_2 - \frac{\langle \mathbf{v}_2, \mathbf{w}_1 \rangle}{\langle \mathbf{w}_1, \mathbf{w}_1 \rangle} \mathbf{w}_1 = \mathbf{v}_2 = (0, 2, 2, 0) \text{ as } \mathbf{v}_2 \perp \mathbf{v}_1.$$

$$\begin{aligned} \mathbf{w}_3 &= (\mathbf{z} - \mathbf{x}_0) - \frac{\langle \mathbf{z} - \mathbf{x}_0, \mathbf{w}_1 \rangle}{\langle \mathbf{w}_1, \mathbf{w}_1 \rangle} \mathbf{w}_1 - \frac{\langle \mathbf{z} - \mathbf{x}_0, \mathbf{w}_2 \rangle}{\langle \mathbf{w}_2, \mathbf{w}_2 \rangle} \mathbf{w}_2 \\ &= (-1, 0, 1, 0) - \frac{0}{4}(1, -1, 1, -1) - \frac{2}{8}(0, 2, 2, 0) \\ &= (-1, -1/2, 1/2, 0). \end{aligned}$$

$$|\mathbf{w}_3| = \left| \left(-1, -\frac{1}{2}, \frac{1}{2}, 0 \right) \right| = \frac{1}{2} |(-2, -1, 1, 0)| = \frac{\sqrt{6}}{2} = \sqrt{\frac{3}{2}}.$$

Modifications of the Gram-Schmidt process

The first modification combines orthogonalization with normalization. Suppose $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ is a basis for an inner product space V . Let

$$\mathbf{v}_1 = \mathbf{x}_1, \quad \mathbf{w}_1 = \frac{\mathbf{v}_1}{\|\mathbf{v}_1\|},$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \langle \mathbf{x}_2, \mathbf{w}_1 \rangle \mathbf{w}_1, \quad \mathbf{w}_2 = \frac{\mathbf{v}_2}{\|\mathbf{v}_2\|},$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \langle \mathbf{x}_3, \mathbf{w}_1 \rangle \mathbf{w}_1 - \langle \mathbf{x}_3, \mathbf{w}_2 \rangle \mathbf{w}_2, \quad \mathbf{w}_3 = \frac{\mathbf{v}_3}{\|\mathbf{v}_3\|},$$

.....

$$\mathbf{v}_n = \mathbf{x}_n - \langle \mathbf{x}_n, \mathbf{w}_1 \rangle \mathbf{w}_1 - \dots - \langle \mathbf{x}_n, \mathbf{w}_{n-1} \rangle \mathbf{w}_{n-1},$$

$$\mathbf{w}_n = \frac{\mathbf{v}_n}{\|\mathbf{v}_n\|}.$$

Then $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n$ is an orthonormal basis for V .

Modifications of the Gram-Schmidt process

Another modification is a recursive process which is more stable to roundoff errors than the original process. Suppose $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ is a basis for an inner product space V . Let

$$\mathbf{w}_1 = \frac{\mathbf{x}_1}{\|\mathbf{x}_1\|},$$

$$\mathbf{v}_2 = \mathbf{x}_2 - \langle \mathbf{x}_2, \mathbf{w}_1 \rangle \mathbf{w}_1,$$

$$\mathbf{v}_3 = \mathbf{x}_3 - \langle \mathbf{x}_3, \mathbf{w}_1 \rangle \mathbf{w}_1,$$

.....

$$\mathbf{v}_n = \mathbf{x}_n - \langle \mathbf{x}_n, \mathbf{w}_1 \rangle \mathbf{w}_1.$$

Then $\mathbf{w}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is a basis for V , $\|\mathbf{w}_1\| = 1$, and \mathbf{w}_1 is orthogonal to $\mathbf{v}_2, \dots, \mathbf{v}_n$. Now repeat the process with vectors $\mathbf{v}_2, \dots, \mathbf{v}_n$, and so on.