

Dilution Problems

$V \uparrow M \downarrow$

400 mL
1 M

$$M_1 V_1 = M_2 V_2$$

2 M

200 mL

Dilution and concentration
Lab 7

Prepared by
Lect. Nora Zawar

Assist. Lec. Zahraa Amer

$$= M_3 V_3$$

$V \downarrow M \uparrow$



Dilution and Concentration

- **Dilution** means when a given solution of a mixture of high concentration is diluted by addition of the suitable diluents or admixture with solution of lower concentration.
- While **concentration** means when a given solution of a mixture of low concentration are concentrated either by addition of active ingredient or by admixture with higher strength solution or by evaporation of the diluents.
- We have different types of dilution either of liquids or solids.




before


after adding solvent

Dilution

$$c_1 \cdot V_1 = c_2 \cdot V_2$$



higher concentration




lower concentration



Dilution Law

- $C_1 V_1 = C_2 V_2$
- The concentration is expressed either by normality, molarity or percent (%).
- Normality is an expression of the concentration of the solution in terms of equivalent per liter of solution (number of gram eq.wt per 1000ml).
- Molarity is the concentration of the solution in terms of moles per liter.



A hand is pouring a clear liquid from a small beaker into a larger Erlenmeyer flask. The flask already contains some liquid, and the new liquid is being added to it. In the background, there are other glassware items like a round-bottom flask and a graduated cylinder, all containing liquid. The entire scene is set against a blue gradient background.

Addition of extra solvent to the existing solution, in order to reduce its concentration is called 'dilution'.



NOTES :

We have two rules wherever they may be applied will simplify the calculation :

- When ratio strength are given, convert them to % before setting.

Ex: $1:10=10\%$

- Wherever proportional parts enter into calculation, reduce them to the lowest terms.
- Ex: 75:25 simplify to 3:1



Dilution of alcohol

- When alcohol is diluted with water a noticeable contraction in volume occurs so it is difficult to calculate the amount of water to be add because alcohol interaction with water by bounding (H-bond) and lead to contraction but this contraction of volume not affect the weight of alcohol and water added.

- Examples

Rx	Boric acid	10gm
	Alcohol 70%	30ml
	Alcohol available	90%

- How many mls of 20% alcohol can be used to prepare 25ml of 10% alcohol?

$$C_1V_1=C_2V_2$$

$$20\% \times V_1 = 10\% \times 25\text{ml}$$

$V_1 = 12.5\text{ml}$ of 20% alcohol and complete the volume to 25ml.



Problems

- If 500ml of 15% v/v solution are diluted to 1500ml. What will be the percentage strength?

$$C_1V_1 = C_2V_2$$

$$15\% \times 500\text{ml} = C_2 \times 1500\text{ml}$$

$$C_2 = 5\%$$

- How many mls of a 1:5000 (w/v) solution of potassium permanganate can be made from 50 ml of a 5% solution?

$$1:5000 = 0.02\%$$

$$C_1V_1 = C_2V_2$$

$$50\text{ml} \times 5\% = 0.02\% \times V_2$$

$$V_2 = 1250\text{ml}$$



Problems

- How much water should be mixed with 5000ml of 85% alcohol to make 50% (v/v) solution?

$$C_1V_1 = C_2V_2$$

$$5000\text{ml} \times 85\% = 50\% \times V_2$$

$$V_2 = 8500\text{ml}$$

$$8500 - 5000 = 3500\text{ml of H}_2\text{O}.$$

- Note
- Standard solution :is a solution of known concentration (normality, molarity and molality) or it's concentration is exactly measured.
- Standardization :is determination of the molarity or normality of the solution.



Reducing and Enlarging Formula

- Pharmacist may have to **reduce or enlarge** the formula in pharmaceutical preparation. In large manufacturing the official formula must be enlarged, while in the pharmacy or on small products the official formula must be reduced



- Factor= desired amount/ specified amount

Examples:

Rx	Codeine phosphate	gr V
	Amaranth solution	℥ XV
	Alcohol 10 %	f3 ss
	D.W. q.s	f3 I

1. Mitt f3 II

2. Mitt f3 ss

Calculation(1):

$5/15 = 0.3$ g of codeine phosphate

$15/15 = 1$ ml of amaranth

f3 ss= 2 ml f3 = 30 ml f 3 II= 60 ml

factor= $60/30 = 2$

$0.3 \times 2 = 0.6$ g of codeine phosphate

$1 \times 2 = 2$ ml of amaranth

$2 \times 2 = 4$ ml of alcohol

$60 \times 3/4 = 45$ ml

$45 - (4 + 2) = 39$ ml



Procedure(1):

- 1. Weigh 0.6 g of codeine phosphate and put it in a beaker.
- 2. Dissolve the amount of codeine phosphate in 39 ml of D.W.
- 3. Add 2 ml of amaranth and 4 ml of alcohol into the content of the beaker.
- 4. Transfer the content of the beaker into a measuring cylinder and complete the volume to 60 ml by D.W.
- 5. Convert the content of the measuring cylinder into a wide mouth bottle and put a suitable label.



Calculation

- $f_{ss} = 15 \text{ ml}$ factor = $15 / 30 = 0.5$
- $0.3 \times 0.5 = 0.15 \text{ g}$ of codeine phosphate
- $1 \times 0.5 = 0.5 \text{ ml}$ of amaranth
- $2 \times 0.5 = 1 \text{ ml}$ of alcohol
- $30 \times 0.5 = 15 \text{ ml}$
- $15 \times 3/4 = 11.25 \text{ ml}$
- $11.25 - (0.5 + 1) = 9.75 \text{ ml}$
- Procedure

Follow the same of the above procedure.



Thank
you!!

